

**WARNING:**

This product must not be applied at a rate greater than that shown on the label.

**FIRST AID INSTRUCTIONS:**

If poisoning occurs, contact a doctor or Poisons Information Centre. If swallowed, and if more than 15 minutes from a hospital, induce vomiting, preferably using Ipecac Syrup APF. If skin contact occurs, remove contaminated clothing and wash thoroughly.

**CONDITIONS OF SALE:**

The manufacturer and distributor warrants that this product contains the active constituents and lignin compounds as specified and is fit for the purpose of correcting nutrient deficiencies when used in accordance with directions under normal conditions of use. No responsibility is accepted by the manufacturer or distributor for use of this product under those conditions which are beyond the control of the manufacturer and distributor.

Manufactured in Australia for:

**TNN INDUSTRIES** pty. ltd.

**1114 JOHNSON ROAD,  
STANHOPE VIC 3623**

**TEL: 03 5857 2422**

**FAX: 03 5857 2350**

**SHAKE WELL  
BEFORE USE**

KEEP OUT OF REACH OF CHILDREN  
**FOR ANIMAL TREATMENT ONLY**



**MINERAL-PLUS**  
**Trace Element Drench for Sheep and Cattle**  
Vitamin A, D, & E

Active constituents:

Cobalt (Co)	as Cobalt lignosulphonate	2.0g
Selenium (Se)	as Selenium: methionine, lignocystine	2.1g
Sulphur (S)	as propylene-phenyl sulphonate	26.7g
Copper (Cu)	as Copper: glycinate, lignosulphonate	3.6g
Zinc (Zn)	as Zinc: methionine, lignosulphonate	7.5g
Manganese (Mn)	as Manganese: cystine, lignosulphonate	5.1g
Iodine (I)	as (+) 3,5 Diiodotyrosine	0.12g
Magnesium (Mg)	as Magnesium: leucine, lignosulphonate	4.5g
Reducing Sugars	as Glucose, Arabinose, Mannose	42.0g
Crude Protein Equivalent	as Amino acids and compounds	305.0g
Propylene Glycol	as 1,2-Propaneolol	100.0g
Micro Fine Kelp		110.0g
Vitamin A	as $\beta$ -carotene	1.125g
Vitamin D	as $\gamma$ -calciferol	7.5mg
Vitamin E	as dl-a-tocopheryl acetate	10.0g

Each litre contains

A supplement for sheep and cattle; to increase dietary intake of certain trace minerals where intake by livestock may be low.

**TNN Mineral Plus employs a unique chelation process to assist in mineral uptake.**

Batch No.: MPD-052    Manf. Date: 01/07/03    Exp. Date: 01/07/07

**DIRECTIONS FOR USE:** Dose Rates for **TNN Mineral Plus**

Lambs and Hoggatts	5 ml	Do not administer to animals more than once in every six weeks or as directed by a Veterinary Surgeon.
Adult Sheep	10 ml	
Rams	12 ml	

Dairy & Beef Cattle (Liveweight kg)	Dosage Rate (ml)	Frequency (weeks)	Dairy & Beef Cattle (Liveweight kg)	Dosage Rate (ml)	Frequency (weeks)
At 1 month	5	2	251 - 300	30	4
50 - 100	10	2	301 - 350	35	5
101 - 150	15	3	351 - 400	40	6
151 - 200	20	3	401 - 450	45	6
201 - 250	25	4	451 kg and over	50	6

Refer back to brochure for feed and water additive dose rates.

- Ensure contents are well mixed by shaking container prior to use.
- Use correct drenching techniques and avoid injury to livestock by maintaining a smooth drenching nozzle.
- Wash drenching equipment thoroughly in clean water, immediately after use.

**WITHHOLDING PERIOD:**

**DO NOT ADMINISTER LATER THAN 7 DAYS BEFORE SLAUGHTER FOR HUMAN CONSUMPTION.**

**SAFETY DIRECTIONS:**

This product may irritate the eyes and skin. Avoid contact with eyes and skin. Wash hands after use. Avoid contamination of food, feed and domestic water supplies.

**FIRST AID INSTRUCTIONS:**

If poisoning occurs, contact a doctor or Poisons Information Centre. If swallowed, and if more than 15 minutes from a hospital, induce vomiting, preferably using Ipecac Syrup APF. If skin contact occurs, remove contaminated clothing and wash thoroughly.

**DISPOSAL:**

Triple or preferably pressure rinse containers before disposal, and dispose of rinsate in a disposal pit away from desirable plants and their roots, and water courses. On-site disposal of undiluted chemicals is unacceptable.

**STORAGE INSTRUCTIONS:**

Store below 30C (room temperature) in original container, in a dry place. Protect from light. Keep container tightly closed.